

Pancyclicity in Claw-free Graphs

R.J. Gould, F. Pfender

*Emory University, Department of Mathematics and Computer Science, Atlanta,
GA 30322, USA*

Abstract

In this paper, we present several conditions for $K_{1,3}$ -free graphs, which guarantee the graph is subpancyclic. In particular, we show that every $K_{1,3}$ -free graph with minimum degree sum $\delta_2 > 2\sqrt{3n+1} - 4$; every $\{K_{1,3}, P_7\}$ -free graph with $\delta_2 \geq 9$; every $\{K_{1,3}, Z_4\}$ -free graph with $\delta_2 \geq 9$; and every $K_{1,3}$ -free graph with maximum degree Δ , $\text{diam}(G) < \frac{\Delta+6}{4}$ and $\delta_2 \geq 9$ is subpancyclic.

Key words: claw-free, pancyclicity, forbidden subgraphs

1 Introduction

If not specified otherwise, we will use notation from [1]. We consider finite simple graphs only. A graph on n vertices is called *subpancyclic* if it contains cycles of every length l with $3 \leq l \leq c(G)$, where $c(G)$ denotes the circumference of G . If G is subpancyclic and hamiltonian, it is called *pancyclic*.

We will always denote the edge set of the graph G by E , and V will denote its vertex set. For some graph H , a graph is said to be *H -free*, if it does not contain an induced copy of H . The complete bipartite graph $K_{1,3}$ is also called the *claw*. The graph Z_4 is a triangle with a path of length four attached to one of its vertices, the graph P_7 is the path on seven vertices.

The degree of a vertex v is denoted by $d(v)$. We will write $\Delta(G)$ or (if no confusion arises) Δ for the maximum degree in G , and $\delta(G)$ or δ for the minimum degree in G . By $\delta_2(G)$ or δ_2 , we will denote the minimum of $\{d(u) + d(v) \mid u, v \in V, uv \notin E\}$.

Email addresses: rg@mathcs.emory.edu (R.J. Gould),
fpfende@mathcs.emory.edu (F. Pfender).

Let C be a cycle in G , and assign some orientation to C . For two vertices $x, y \in V(C)$, the notation xCy will stand for the path from x to y along C following the orientation of C . An xy -path P in G is called a *shortening path of C* , if $V(P) \cap V(C) = \{x, y\}$ and $|P| < \min\{|xCy|, |yCx|\}$. An edge $xy \notin E(C)$ with $x, y \in V(C)$ is called a *chord of C* .

We will start by proving the following Lemma.

Lemma 1 *Let G be a claw-free graph with $\delta_2(G) \geq 9$. Suppose, for some $m > 3$, G has an m -cycle C , but no $(m-1)$ -cycle. Then there is no shortening path of C .*

As we will see, Lemma 1 has several interesting consequences.

Theorem 2 *Let G be a claw-free graph with maximum degree Δ and $\delta_2(G) \geq 9$. If $\text{diam}(G) < \frac{\Delta+6}{4}$, then G is subpancyclic.*

Theorem 3 *Let G be a claw-free graph with minimum degree δ and $\delta_2(G) \geq 9$. If G is not a line graph, and $\text{diam}(G) < \frac{\delta+3}{2}$, then G is subpancyclic.*

Theorem 4 *Let G be a $\{K_{1,3}, Z_4\}$ -free graph with $\delta_2 \geq 9$. Then G is subpancyclic. If G is 2-connected, then G is pancyclic.*

Theorem 5 *Let G be a $\{K_{1,3}, P_7\}$ -free graph with $\delta_2 \geq 9$. Then G is subpancyclic.*

Theorem 6 *Let G be a claw-free graph on $n \geq 5$ vertices with $\delta_2 > 2\sqrt{3n+1}-4$. Then G is subpancyclic.*

From Theorem 6 we obtain as a corollary the following Theorem of Trommel, Veldman and Verschut [2]:

Theorem 7 *Let G be a claw-free graph on $n \geq 5$ vertices. If the minimum degree δ is $\delta > \sqrt{3n+1}-2$, then G is subpancyclic.*

In the proofs of Theorems 2-6, we will frequently use the following theorem from Flandrin, Fournier and Germa [4], and its corollaries:

Theorem 8 *Let G be a claw-free graph. Then the graph $\langle N \rangle$ induced by the neighborhood N of any vertex x falls in one of three cases:*

1. $\langle N \rangle$ is hamiltonian.
2. $\langle N \rangle$ consists of two complete subgraphs G_1 and G_2 , connected with some edges, all of them having a common vertex in G_1 .
3. $\langle N \rangle$ consists of two complete subgraphs with no edges in between.

Corollary 9 *Let G be a claw-free graph with maximum degree Δ . Then G*

contains cycles of length l for all l with $3 \leq l \leq \lceil \Delta/2 \rceil + 1$.

Proof: The proof is obvious. \square

Corollary 10 *Let G be a claw-free graph with minimum degree δ . If G is not a line graph, then G contains cycles of length l for all l with $3 \leq l \leq \delta + 1$.*

Proof: Observe that G is a line graph if the neighborhoods of all vertices are in the third class of Theorem 8. Therefore, there is a vertex x with $\langle N(x) \rangle$ in the first or second class of Theorem 8. In either case, $\langle N(x) \rangle$ is traceable, implying $\langle N(x) \cup \{x\} \rangle$ is pancyclic. \square

2 Proof of Lemma 1

Suppose instead P is a shortest shortening path. We will distinguish two cases.

Case 1 *Suppose P is a chord ($P = xy$).*

Pick two chords u_1u_2 and v_1v_2 , such that $u_1, u_2 \in xCy, v_1, v_2 \in yCx$, where both chords are minimal in the sense that there is no other chord uv with $u, v \in u_1Cu_2$ or $u, v \in v_1Cv_2$. This does not exclude the possibility of one or both of these chords being identical with xy .

Let $K := \{v \in V(C) \mid \exists u \in V(C) : uv \text{ is a chord}\}$, $L := V(C) - K$. If there is a shortening path of C with length exactly two with both its endvertices in u_1Cu_2 (v_1Cv_2), pick such a shortening path $s_1s_2s_3$ ($t_1t_2t_3$), such that s_1Cs_3 (t_1Ct_3) is as short as possible, else set $s_1 = s_2 = u_1, s_3 = u_2$ ($t_1 = t_2 = v_1, t_3 = v_2$).

Let a_1, a_2, \dots, a_r be the vertices of $s_1^+Cs_3^- \cap L$ (in order), let b_1, b_2, \dots, b_l be the vertices of $t_1^+Ct_3^- \cap L$. Without loss of generality, by symmetry we may assume that $l \geq r$. Further, if $l = r$ we may assume that $d(b_i) \geq 5$ for all $1 \leq i \leq l$ (since they belong to L , there are no edges between the a_i and the b_j , so $\delta_2(G) \geq 9$ guarantees the statement).

Now we will construct a cycle $C' \subset \langle C \cup s_2 \rangle$ with $m - r - 1 \leq |C'| \leq m - 1$, which we will then extend to a C_{m-1} to get a contradiction.

Start with the cycle $s_1s_2s_3Cs_1$. Note that $c = |s_1s_2s_3Cs_1| \leq m - 1$. If $c \geq m - r - 1$, this cycle is the desired C' . Otherwise, $s_1^+Cs_3^- \cap K \neq \emptyset$ and we can pick a vertex $u \in s_1^+Cs_3^- \cap K$. Then u has an edge to some vertex $v \in s_3^+Cs_1^-$. There can't be an edge v^-v^+ , else there is a C_{m-1} . There is no claw centered at v , so $v^+u \in E$ or $v^-u \in E$. Therefore u can be inserted in the cycle between v

and one of its neighbors to extend the cycle. If two vertices $u, w \in s_1^+ C s_3^- \cap K$ share the same neighbors $v, v^+ \in s_3^+ C s_1^-$, then all of $u C w$ (or $w C u$) can be inserted between v and v^+ to extend the cycle. Thus, any number of vertices in $s_1^+ C s_3^- \cap K$ can be inserted (we don't have control about the number of vertices of $s_1^+ C s_3^- \cap L$ inserted in the process). With this process, we insert $m - r - 1 - c$ vertices out of $s_1^+ C s_3^- \cap K$. The resulting cycle C' is of the desired length, since at most r vertices out of L were inserted.

To extend C' , consider $b_1, b_4, b_7, \dots, b_{3\lceil l/3 \rceil - 2}$. Since $t_1 C t_3$ is the shortest such segment possible, these vertices have pairwise disjoint neighborhoods. Further, none of them is a neighbor of s_2 , else there is a claw at s_2 . By Theorem 8, C' can be extended through the neighborhoods of these vertices by any number of vertices up to $d(b_i) - 2$ for each $b_i, i = 1, 4, \dots$.

If $l = r$, then $d(b_i) \geq 5$, so this extends C' by up to $3\lceil l/3 \rceil \geq r$ vertices, resulting in a C_{m-1} .

If $3 \leq r < l$, let $d := \min\{4, d(b_1), d(b_4), \dots\}$. Then C' is extendable by $\sum_{i=0}^{\lceil l/3 \rceil - 1} (d(b_{1+3i}) - 2) \geq d - 2 + (\lceil l/3 \rceil - 1)(7 - d) \geq 3\lceil l/3 \rceil - 1$ vertices, yielding a C_{m-1} .

If $1 \leq r < l = 3$, consider b_1 and b_3 . One of them has degree at least 5, so we can extend by up to 3 vertices, which is again enough.

If $r = 1, l = 2$, the only problem would be if $d(b_1) = d(b_2) = 2$, else we could extend by one, which is enough. But then, $d(a_1) \geq 7$, and by a symmetric argument we can find a cycle $C'' \subset \langle C \cup t_2 \rangle$ which includes $a_1^- a_1 a_1^+$, and $m - 3 \leq |C''| \leq m - 1$. This cycle can now be extended around a_1 to a C_{m-1} .

Finally, if $r = 0$, C' is already a C_{m-1} . This contradiction concludes the argument, hence C has no chords.

Case 2 Suppose P has length ≥ 2 ($P = z_0 z_1 z_2 \dots z_l$, with $x = z_0, y = z_l$).

Assume, that P is chosen such that $k = |x C y|$ is minimal. Observe, that $k > l - 1$ (else $k = l - 1$, and a C_{m-1} is easily found). Let $v_0 = y, v_1 = y^+, \dots, v_{m-k+1} = x$. Since k is minimal, $x^+ z_1 \notin E$. Since C is chordless, $x^+ v_{m-k} \notin E$. Thus $v_{m-k} z_1 \in E$ to prevent a claw at x . A symmetric argument shows that $v_1 z_{l-1} \in E$. Now $m - k \geq k$, else k would not have been minimal.

Consider $C' = x P y C x$. We know that $m - k + l + 1 = |C'| \leq m - 2$. We will now extend C' to a C_{m-1} to get the contradiction. None of the edges $v_i z_j, 2 \leq i \leq m - k - 1, 0 \leq j \leq l$ exists, else let j be minimal, such that for some $2 \leq i \leq m - k - 1$, there is an edge $v_i z_j$ ($j \geq 1$, else chord). To prevent a claw at z_j , $z_{j+1} v_i \in E$ is necessary. But now, consider the paths $P' = v_i z_{j+1} P y$ and $P'' = v_i z_j P x$. Both of them are shorter than P . Since P is the shortest shortening path, P' and P'' can not be shortening paths, thus

$1 + l - j = |P'| \geq |yCv_i| = i + 1$, and $j + 2 = |P''| \geq |v_iCx| = m - k - i + 2$. But this implies that $l \geq m - k \geq k$, a contradiction to P being a shortening path.

Now, note that none of the neighborhoods of $v_2, v_5, \dots, v_{3\lfloor k/3 \rfloor - 1}$ intersect, else k was not minimal.

If $k \geq 6$, let $d := \min\{4, d(v_2), d(v_5), \dots, d(v_{3\lfloor k/3 \rfloor - 1})\}$. We can extend C' around $v_2, v_5, \dots, v_{3\lfloor k/3 \rfloor - 1}$ by up to $\sum_{i=0}^{\lfloor k/3 \rfloor - 1} (d(v_{2+3i}) - 2) \geq d - 2 + (\lfloor l/3 \rfloor - 1)(7 - d) \geq 3\lfloor k/3 \rfloor - 1 \geq k - 3$ vertices to get a C_{m-1} , just like in the first case.

If $k = 5$, $|C'| = m + l - 4 \geq m - 2$. As either $d(v_2) \geq 5$ or $d(v_4) \geq 5$, we can extend around it by one vertex, and we have our contradiction.

If $k \leq 4$, then $m - 2 \geq |C'| \geq m - k + l + 1 \geq m - 4 + 2 + 1 = m - 1$, a contradiction. \square

3 Proofs of the Theorems

Proof of Theorem 2: Suppose G is not subpancyclic. Then for some m , G has a C_m , but no C_{m-1} . By Corollary 9, $m \geq \frac{\Delta+6}{2}$. But now the diameter condition guarantees a shortening path, which is impossible by Lemma 1. \square

Proof of Theorem 3: Suppose G is not subpancyclic. Then for some m , G has a C_m , but no C_{m-1} . By Corollary 10, $m \geq \delta + 3$. But now the diameter condition guarantees a shortening path, which is impossible by Lemma 1. \square

Proof of Theorem 4: Suppose G is not subpancyclic. Then for some m , G has a C_m , but no C_{m-1} . By Corollary 9, $m \geq 6$. By Lemma 1, $C := C_m$ has no chords. By the degree condition, there is a vertex $v \in V(C)$ with $d(v) \geq 5$.

If $m = 6$, the neighborhood of v is split into two complete subgraphs, not connected by edges. Else there was a 5-cycle in $N(v) \cup v$ by Theorem 8 (take a P_4 in $N(v)$, and connect both its ends with v). Without loss of generality, let $x, y \in N(v) \cap N(v^+)$ (so $xv^-, yv^- \notin E, xy \in E$). Observe that $xv^{++}, yv^{++} \notin E$, else there is a 5-cycle. Further x and y can not be adjacent to any other vertex of C , else there is a shortening path of C , which is not possible by Lemma 1. Now yxv^+Cv^- form a Z_4 , a contradiction.

If $m \geq 7$, there exist $z \in V - V(C), y \in V(C)$, such that $z \in (N(y) \cap N(y^+)) - N(y^{++})$. But then zyy^+Cy^{5+} form a Z_4 (Again, z can not be adjacent to any of y^{3+}, y^{4+}, y^{5+} , else there was a shortening path of C).

If G is 2-connected, then it is hamiltonian by a result of Brousek, Ryjáček and Favaron [3], thus G is pancyclic. \square

Proof of Theorem 5: Suppose G is not subpancyclic. Then for some m , G has a C_m , but no C_{m-1} . By Corollary 9, $m \geq 6$. By Lemma 1, $C := C_m$ has no chords.

If $m = 6$, let $v \in V(C)$ be the vertex on C with the largest degree. The degree condition guarantees $d(v) \geq 5$. By Theorem 8, the neighborhood of v is split into two complete subgraphs of size at most 3, not connected by edges, else there is a C_5 in $N(v) \cup v$. Hence, $|N(v) \cap N(v^-)|, |N(v) \cap N(v^+)| \in \{1, 2\}$, with one of them being 1 only in the case that $d(v) = 5$. Let $x \in N(v) \cap N(v^+), y \in N(v) \cap N(v^-)$. Clearly $xy \notin E$ or a C_5 is immediate. Now neither of the two edges xv^{++}, yv^{--} can exist by the following argument: If $|N(v) \cap N(v^+)| = 2$, then xv^{++} completes a 5-cycle. If $|N(v) \cap N(v^+)| = 1$, then $d(v) = 5$, and therefore $d(z) \geq 4$ for all $z \in V(C)$ (a z with a smaller degree would guarantee a vertex of degree ≥ 6 in the chordless C , contradicting $d(v)$'s maximality). In particular $d(v^+) \geq 4$. Let $x' \in N(v^+) - \{v, v^{++}, x\}$. Then $x'v^{++} \in E$ to prevent a claw at v^+ . If $xv^{++} \in E$, then $xv^{++}x'v^+vx$ is a C_5 . Hence, in either case $xv^{++} \notin E$. The argument against $yv^{--} \in E$ is symmetric.

Further, x, y can not have any other adjacencies on C , else a shortening path of C exists, a contradiction to Lemma 1. Now xv^+Cv^-y is a P_7 , a contradiction.

If $m = 7$, observe that there are at most two vertices on C with degree ≤ 4 . Thus there is a vertex $v \in V(C)$ with $d(v), d(v^+), d(v^-) \geq 5$. By Theorem 8, the neighborhood of v is split into two complete subgraphs, not connected by edges, else there is a C_6 in $N(v) \cup v$. Without loss of generality, let $x, y \in N(v) \cap N(v^+)$. Then $xv^{++} \notin E$, else there is a C_6 in $v^+ \cup N(v^+)$, using v, v^+, v^{++}, x, y and one other neighbor of v^+ . Further, x has no other neighbors on C , else a shortening path of C exists. But now xv^+Cv^- is a P_7 , a contradiction.

If $m \geq 8$, vCv^{6+} forms a P_7 for any $v \in V(C)$, a contradiction.

Hence, G is subpancyclic. \square

Proof of Theorem 6: Suppose G is not subpancyclic. Then for some m , G has a C_m , but no C_{m-1} . By Corollary 9, $m \geq \Delta/2 + 3 \geq \delta_2/4 + 3$.

Case 1 Suppose $n < 12$.

Note that the degree sum condition implies the following bounds on δ_2 :

$$n \in \{6, 7, 8\} \Rightarrow \delta_2 \geq n - 1,$$

$$n \in \{9, 10\} \Rightarrow \delta_2 \geq n - 2,$$

$$n = 11 \Rightarrow \delta_2 \geq n - 3.$$

Consider all the possible values for m . Since $n \geq 5$, $m \geq \frac{\Delta}{2} + 3 > \frac{\sqrt{3n+1}}{2} + 3 \geq 5$. Say, $C = C_m = v_1 \dots v_m v_1$.

If $m = 6$, then the only chords C could have are of the form $v_i v_{i+3}$. But then claw-freeness forces either $v_i v_{i+2}$ or $v_i v_{i+4}$, which leads to a C_5 . So C has no chords.

For $n \leq 8$, there are at least

$$\sum_{i=1}^6 (d(v_i) - 2) \geq 3\delta_2 - 12 \geq 3(n-1) - 12 = 3n - 15$$

edges from C to $V - C$, but at most $3(n-6) = 3n - 18$ edges from $V - C$ to C , since no vertex in $V - C$ can have more than three neighbors on C without producing a C_5 . Thus, $n \geq 9$.

For $9 \leq n \leq 10$, the same count shows that there are exactly $3n - 18$ edges from C to $V - C$, hence every vertex of $V - C$ has exactly three neighbors on C . To avoid a claw and a C_5 , all three have to be in a row. If two of the vertices $u, w \in V - C$ are adjacent, a C_5 can easily be found. But now $d(u) + d(w) = 6 < \delta_2$, a contradiction.

For $n = 11$, there are at least $3\delta_2 - 12 \geq 12$ edges from C to $V - C$, so out of the five vertices in $V - C$, at least two vertices $u, w \in V - C$ have three neighbors on the cycle, and two more vertices $x, y \in V - C$ have at least two neighbors on the cycle. If any of the edges uw, ux, uy, wx, wy exists, a C_5 can easily be found. Since $\delta_2 \geq 8$, both u and w must be adjacent to the remaining vertex z . But now again, a C_5 can be found.

If $m = 7$, the only possible chords are of the form $v_i v_{(i+3) \bmod 7}$. To avoid claws, all chords of this form have to exist if one exists. But now $v_1 v_2 v_5 v_6 v_7 v_4 v_1$ is a C_6 . Therefore, C has no chords. This yields immediately $n \geq 8$. Observe, that for $n < 12$, the degree sum condition ensures that $\delta_2 \geq n - 3$. Now a similar count as in the last case gives at least

$$\sum_{i=1}^7 (d(v_i) - 2) \geq \frac{7}{2}\delta_2 - 14 \geq \frac{7}{2}(n-3) - 14 = 3n - 21 + \frac{n-7}{2}$$

edges going out of C , with at most $3(n-7)$ going in, a contradiction.

If $m = 8$ and C has a chord, then C has exactly the chords (after a cyclic renumbering of the vertices) $v_1 v_5, v_1 v_6, v_2 v_5, v_2 v_6$. If any of those are missing, there is a claw, if there are any more than those, there is a C_7 . But now the degree sum condition forces v_3 or v_8 to have a neighbor outside the cycle, say $v_3 x \in E$. To avoid a claw, $v_2 x \in E$ or $v_4 x \in E$. But this again yields a C_7 . So

C has no chords, and a similar count as before yields

$$\sum_{i=1}^8 (d(v_i) - 2) \geq 4\delta_2 - 16 \geq 4(n - 3) - 16 = 4n - 28 > 3(n - 8),$$

a contradiction.

If $m = 9$, a similar count shows the existence of chords. But if there is a chord, claw-freeness forces the appearance of a K_4 of the form $v_i v_{i+1} v_{i+4} v_{i+5}$ inside $\langle C \rangle$, say at $v_1 v_2 v_5 v_6$.

Now v_8 has no neighbors outside C : Suppose $x \in V - C$, $xv_8 \in E$. To prevent a claw at v_8 , x has to be adjacent to v_7 or v_9 . But then the 7-cycle $C' = v_2 v_5 C v_2$ can be extended to a C_8 through x .

If $v_8 v_3 \in E$, then v_3 is adjacent to either v_7 or v_9 to avoid a claw at v_8 . But then again, C' can be extended through v_3 . The symmetric argument shows that $v_8 v_4 \notin E$. Further, if $v_8 v_2 \in E$, then $v_8 v_3 \in E$ to prevent a claw at v_2 , which is not possible. The symmetric argument shows that $v_8 v_5 \notin E$. So $d(v_8) = 2$. But this implies that $d(v_3) \geq n - 5$. We know that v_3 is not adjacent to v_8 , v_1 and v_5 . Further, v_3 can not be adjacent to v_9 without creating a claw at v_9 . Thus, v_3 is adjacent to all other vertices, in particular $v_3 v_6, v_3 v_7 \in E$. But now, C' can be extended through v_3 , a contradiction.

If $m \geq 10$, a chord is guaranteed, again. Consider a chord $v_i v_j$, such that $|v_i C v_j|$ is minimal. Now find a chord $v_r v_s$ on $v_j C v_i$, such that there is no other chord within $v_r C v_s$. Either all vertices in $v_r C v_s$ or all vertices in $V_i C v_j$ have chords, since there is at most one vertex outside C , and all vertices with degree at most 3 have to be pairwise adjacent. Say all vertices in $v_r C v_s$ have chords. Now, similar to the first case in the proof of Lemma 1, insert all but one of $v_{r+1} C v_{s-1}$ into $v_s C v_r v_s$ to construct a C_{m-1} .

Case 2 Suppose $n \geq 12$, $m \geq \delta_2/2 + 3$.

By Lemma 1, C has no chords ($n \geq 12$ guarantees $\delta_2 \geq 9$). Thus there are

$$\sum_{v \in V(C)} (d(v) - 2) \geq m \left(\frac{\delta_2}{2} - 2 \right)$$

edges from C to $G - C$. On the other hand, every vertex in $G - C$ can have at most three neighbors on C , otherwise C has a shortening path, which is impossible by Lemma 1. So

$$m \left(\frac{\delta_2}{2} - 2 \right) \leq 3(n - m),$$

thus

$$3n \geq m\left(\frac{\delta_2}{2} + 1\right) \geq \left(\frac{\delta_2}{2} + 3\right)\left(\frac{\delta_2}{2} + 1\right) > (\sqrt{3n+1} + 1)(\sqrt{3n+1} - 1) = 3n,$$

a contradiction.

Case 3 Suppose $n \geq 12$, $m < \delta_2/2 + 3$.

Let $d := \lceil \delta_2/2 \rceil$, so $m \leq d + 2$. By Corollary 9, we know that $m \geq \Delta/2 + 3 \geq d/2 + 3$, particularly $m \geq 6$. By Lemma 1, C has no chords. Let $C = v_1v_2 \dots v_mv_1$. Since all vertices of degree $< d$ have to be pairwise adjacent, we may assume that $d(v_i) \geq d$ for $3 \leq i \leq m$. For $i = 1, 2, \dots, m-1$, let $N_i := N(v_i) \cap N(v_{i+1})$, let $N_m := N(v_m) \cap N(v_1)$. Since G is claw-free, every vertex adjacent to C lies in some N_i . Note, that if $d(v_i) \geq d$, then $N_{i-1} \cap N_i = \emptyset$, and N_{i-1} and N_i induce complete subgraphs, otherwise, $\langle N(v_i) \rangle$ is traceable by Theorem 8, so we can find cycles of any length up to $d(v_i) + 1$ in $\langle N(v_i) \cup v_i \rangle$, in particular one of length $m - 1$.

Now we claim that there can not be any edges or 2-paths between N_i and N_j , for $3 \leq i < j \leq m - 1$. If $j - i \geq 4$, an edge or 2-path leads to a shortening path of C , a contradiction to Lemma 1. If $j - i \leq 3$ and $m \geq 7$, one can easily find a cycle of length at most 6 through that edge or 2-path, v_{i+1} and v_j , which we can then extend to a C_{m-1} , using any number of vertices out of $N(v_j)$. If $m = 6$, then $j - i \leq 2$, and one can easily find a cycle of length at most 5 through that edge or 2-path, v_{i+1} and v_j , which we can then extend to a C_5 , using any number of vertices out of $N(v_j)$.

Since all vertices of degree less than d have to be pairwise adjacent, we can now guarantee, after possibly renumbering the vertices of C , that all such vertices in $H := \bigcup N_i \cup C$ must lie in $N_m \cup N_1 \cup N_2 \cup \{v_1, v_2\}$.

Our next claim is, that for two vertices $x, y \in N_i, 3 \leq i \leq m - 1$, their neighborhoods intersect as follows: $N(x) \cap N(y) = N_i \cup \{v_i, v_{i+1}\} - \{x, y\}$. We already established that it is at least of that size, since $\langle N_i \rangle$ is complete. But it can not be bigger; for suppose, there is a $z \in (N(x) \cap N(y)) - H$. Then, z is not adjacent to v_i . Therefore, the neighborhood of x is traceable by Theorem 8, and since $d(x) \geq d$, we can find a C_{m-1} in $\langle N(x) \cup x \rangle$.

Let $M_i := \{z \in V - H : zx \in E \text{ for some } x \in N_i\}$. Since $|N_i| \leq m - 4$ for all $3 \leq i \leq m - 1$ (else you can find a C_{m-1} in $\langle N_i \cup \{x_i, x_{i+1}\} \rangle$, since $\langle N_i \rangle$ is complete), and the degree of vertices in N_i is at least d , every $x \in N_i$ has at least $d - m + 3$ neighbors outside $N_i \cup \{x_i, x_{i+1}\}$. Thus, $|M_i| \geq (d - m + 3)|N_i|$ for $3 \leq i \leq m - 1$. Further, the M_i are disjoint, otherwise there would be 2-paths between the N_i .

But now we see that

$$\begin{aligned}
n &\geq |C| + |N_m \cup N_1 \cup N_2| + \sum_{i=3}^{m-1} |N_i \cup M_i| \\
&\geq |C| + |N_m \cup N_1 \cup N_2| + \sum_{i=3}^{m-1} (d - m + 4) |N_i| \\
&\geq^* m + \frac{(d(v_m)-2)+(d(v_1)-2)+(d(v_3)-2)+(d-m+4)\sum_{i=4}^{m-1}(d(v_i)-2)}{2} \\
&\geq \frac{d}{2} + 3 + \frac{d-2+\delta_2-4+(d-m+4)(m-4)(d-2)}{2} \\
&\geq^{**} \frac{4d-1+(2d-4)(d-2)}{2} \\
&> d^2 - 2d + 3 \\
&> n,
\end{aligned}$$

where \geq^* results from a count that counts every vertex in the N_i at most twice, and \geq^{**} comes from the fact, that for $d \geq 2$,

$$\min_{d/2+3 \leq m \leq d+2} ((d - m + 4)(m - 4)) = 2d - 4,$$

and $\delta_2 \geq 2d - 1$. This contradiction concludes the proof. \square

4 Sharpness

In this section we demonstrate the sharpness of some of the results.

The following family of graphs (see also figure 1) demonstrates the sharpness of the bound on δ_2 in Lemma 1. Let $k \geq 4$, and let H_1, \dots, H_{2k} be $2k$ disjoint copies of K_5 , and $u_i v_i$ an edge of H_i ($i = 1, \dots, 2k$). Now the graph F_k is obtained from $\bigcup_{i=1}^{2k} H_i - u_i v_i$ by adding the edges $v_1 u_2, v_2 u_3, \dots, v_{2k-1} u_{2k}, v_{2k} u_1$ and the edges $u_1 v_k, u_1 u_{k+1}, u_{2k} v_k, u_{2k} v_{k+1}$. We have $\delta_2(F_k) = 8$, and there is a C_{6k} with chords, but no C_p for $5k + 2 < p < 6k$.

The graph G in figure 2 shows that in Theorem 5 P_7 -free can not be replaced by P_8 -free. This graph is $\{K_{1,3}, P_8\}$ -free with $\delta_2 = 10$, and G contains a C_8 but no C_7 .

The degree bounds in Theorem 6 and Theorem 7 are sharp. Consider the following family of graphs from [2] :

For any integer $p \geq 2$, we define the graph G_p as follows. Let H_1, \dots, H_p be p disjoint copies of K_{3p-2} , and $u_i v_i$ an edge of H_i ($i = 1, \dots, p$). Now G_p is obtained from $\bigcup_{i=1}^p H_i - u_i v_i$ by adding the edges $v_1 u_2, v_2 u_3, \dots, v_{p-1} u_p, v_p u_1$.

The graph G_p is both hamiltonian and claw-free. Furthermore, we have $\delta(G_p) = 3p - 3$ and $|V(G_p)| = p(3p - 2)$, implying that $\delta(G_p) = \sqrt{3n + 1} - 2$. It is

obvious that G_p does not contain C_{3p-1} and hence G_p is not (sub)pancyclic.

References

- [1] B. Bollobás, **Extremal Graph Theory**, Academic Press Inc, London, 1978.
- [2] H. Trommel, H.J. Veldman, A. Verschut, **Pancyclicity of claw-free hamiltonian graphs**, Discrete Mathematics 197/198 (1999) 781-789.
- [3] J. Brousek, Z. Ryjáček, O. Favaron, **Forbidden subgraphs, hamiltonicity and closure in claw-free graphs**, Discrete Mathematics 196 (1999) 29-50.
- [4] E. Flandrin, I. Fournier, A. Germa, **Pancyclism in $K_{1,3}$ -free graphs**, Ph.D Thesis, 1990.

1 1, 4 3 1




Fig. 1. Graph F_k




Fig. 2. Graph G