Primes in short arithmetic progressions

MSC-index 11N35, 11N13

Keywords: Large Sieve, Selberg's Sieve

The Large Sieve inequality in the form

$$\sum_{q\le Q} q \sum_{a=1}^{q} \left| \sum_{\substack{n\le N \\ n\equiv a \pmod q}} a_n - \frac{1}{q}\sum_{n\le N} a_n \right|^2 < (N+Q^2)\sum_{n\le N}|a_n|^2$$

is essentially optimal. However, in several applications many of the a_n vanish, and one might expect better estimates then. In fact, such estimates were given by P. D. T. A. Elliott[1]. He showed the following estimate:

Theorem 1 *N and Q be integers, a_p be complex numbers for all primes $p \le N$. Then we have the estimate*

$$\sum_{q\le Q}(q-1)\sum_{(a,q)=1}\left| \sum_{\substack{p\le N \\ p\equiv a \pmod q}} a_p - \frac{q}{\varphi(q)}\sum_{p\le N} a_p \right|^2 \ll_\epsilon \left(\frac{N}{\log N} + Q^{54/11+\epsilon}\right)\sum_{p\le N}|a_p|^2$$

Under GRH, $Q^{54/11}$ may be replaced by $Q^{4+\epsilon}$. In analogy to the large sieve, he conjectured that one may replace this term by $Q^{2+\epsilon}$.

Using a completely different approach, Y. Motohashi[4] showed that

$$\sum_{q\le Q}\sum_{\chi \pmod q}{}^{*}|\pi(x,\chi)|^2 \le \frac{(2+o(1))x^2}{\log x \log x/Q^{1/2}} \tag{1}$$

for $x > Q^{5+\epsilon}$, where $\pi(x,\chi) = \sum_{p\le x}\chi(p)$. He also conjectured, that $Q^{5+\epsilon}$ may be replaced by $Q^{2+\epsilon}$.

Here we will combine the Large Sieve principle with Selberg's sieve to prove the conjecture of Elliott and give a version of (1) valid for $x > Q^{2+\epsilon}$.

I would like to thank D. R. Heath-Brown for his help on Proposition 9 which allowed me to reduce the exponent to $2+\epsilon$, and the referee for pointing out some mistakes.

Theorem 2 *Let N and Q be integers with $N > Q^{2+\epsilon}$, a_p be complex numbers for any prime $p \le N$, and let $2 \le R \le \sqrt{N}$ be an integer. Then we have the estimate*

$$\sum_{q\le Q}\sum_{\chi \pmod q}{}^{*}\left|\sum_{p\le N} a_p\chi(p)\right|^2 \ll_\epsilon \frac{N}{\log N}\sum_{p\le N}|a_p|^2$$

As this estimate is the analogue of the large sieve estimate, we can give analogues of Halász-type inequalities, too. As there is a variety of different large value estimates, the same is true for these bounds. However, since the optimal estimate depends on the particular application, we only mention the following:

Theorem 3 *Let q be an integer. Let $\mathcal{C}$ be a set of characters $\pmod q$, a_p be complex numbers for any prime $p \le N$. Then we have for $k = 2, 3$ or,*

if q is cubefree, for any integer $k \geq 2$, the estimates

$$\sum_{\chi \in \mathcal{C}} \left| \sum_{p \leq N} a_p \chi(p) \right|^2 \leq \left(\frac{N}{\log R} + c_{k,\epsilon} N^{1-1/k} q^{(k+1)/(4k^2)+\epsilon} |\mathcal{C}| R^{2/k} \right) \sum_{p \leq N} |a_p|^2$$

and

$$\sum_{\chi \in \mathcal{C}} \left| \sum_{p \leq N} a_p \chi(p) \right|^2 \leq \left(\frac{N}{\log R} + R^2 |\mathcal{C}| \sqrt{q} \log q \right) \sum_{p \leq N} |a_p|^2.$$

If $\mathcal{C}$ is a set of characters to moduli $q \leq Q$, the same bounds apply with q replaced by Q^2, where k can be chosen arbitrarily, if all occuring values of q are cubefree, and $k = 2, 3$ otherwise.

From this we conclude immediately

Corollary 4 *We have for $x > Q^{2+\epsilon}$ the estimate*

$$\sum_{q \leq Q} \sideset{}{^*}\sum_{\chi \pmod q} |\pi(x, \chi)|^2 \leq C_\epsilon \frac{x^2}{\log^2 x}$$

Moreover, for $x > Q^{3+\epsilon}$ this can be made completely explicit:

$$\sum_{q \leq Q} \sideset{}{^*}\sum_{\chi \pmod q} |\pi(x, \chi)|^2 \leq \frac{(2 + o(1))x^2}{\log x \log x / Q^3}$$

We can also consider a single character:

Corollary 5 *Let χ be a complex character. Then we have*

$$|\pi(x, \chi)| \leq \left(\left(\frac{1 + \phi/\alpha}{2 - 2\phi/\alpha} \right)^{1/2} + o(1) \right) \frac{x}{\log x},$$

where $\alpha = \frac{\log x}{\log q}$ and $\phi = \frac{1}{4}$ if q is cubefree, and $\phi = \frac{1}{3}$ otherwise.

Note that this estimate is nontrivial as soon as $x > q^{3/4}$ resp. $x > q$, depending on whether q is cubefree or not. With a little more work, we obtain the following statement.

Corollary 6 *Let D, x, Q be parameters with $x > Q^{1+\epsilon}D^2$. Let N be the number of moduli $q \leq Q$, such that there is some primitive character χ of order $d \leq D$ and some d-th root of unity ζ, such that there is no prime $p \leq x$ with $\chi(p) = \zeta$. Then we have $N \ll_\epsilon D$.*

This was proven by Elliott with $D = 3$ under the condition $x > Q^{54/11+\epsilon}$.

We begin the proof of our Theorems with the following two sieve principles.

Lemma 7 (Bombieri) *Let $V, (\cdot, \cdot)$ be an inner product space, $v_i \in V$. Then for any $\Phi \in V$ we have*

$$\sum_i |(\Phi, v_i)|^2 \leq \|\Phi\|^2 \max_i \sum_j |(v_i, v_j)|$$

This is Lemma 1.5 in [3].

Lemma 8 (Selberg) *Let R, N be integers, such that $R^2 < N$. Then there is a function g, which has the following properties:*

1. *$g(1) = 1$, $|g(n)| \leq 1$ for $n \leq R$, $g(n) = 0$ for $n > R$.*

2. $\sum_{n\leq N}((1*g)(n))^2 \leq \frac{N}{\log R} + R^2$

This is the usual formulation of Selberg's sieve when used to count the set of primes $\leq N$, see e.g. [2], chapter 3, especially Theorem 3.3. In the sequel, we will denote the function given by Lemma 8 with g and set $f = (1*g)^2$. We will have to bound character sums involving f, these computations are summarized in the following Proposition.

Proposition 9 *Let $\chi \pmod q$ be a character, R, N, f and g as in Lemma 8, and define $S = \sum_{n\leq N} f(n)\chi(n)$.*

1. *If χ is principal, we have $|S| < \frac{N}{\log R} + R^2$.*

2. *Assume that χ is nonprincipal. Then we have for any fixed A the estimate $\sum_{\nu=1}^{\infty} f(\nu)\chi(\nu)e^{-\log^2(\nu/N)} \ll_{\epsilon,A} R^2 q^{1/2}\left(\frac{N}{R^2 q}\right)^{-A}$.*

3. *If χ is nonprincipal, we have the bounds $|S| \leq R^2\sqrt{q}\log q$ and $|S| \leq c_{k,\epsilon}R^{2/k}N^{1-1/k}q^{(k+1)/(4k^2)+\epsilon}$ for $k = 2, 3$, or, if q is cubefree, for $k \geq 2$ arbitrary.*

Proof: The first assertion is already contained in Lemma 8.

Assume now that χ is nonprincipal. Then we have

$$\left|\sum_{n\leq N} f(n)\chi(n)\right| = \left|\sum_{n\leq N}\left(\sum_{d|n} g(d)\right)^2 \chi(n)\right|$$

$$
\begin{aligned}
&= \left| \sum_{d_1,d_2 \le R} g(d_1)g(d_2)\chi\left([d_1,d_2]\right) \sum_{n \le N/[d_1,d_2]} \chi(n) \right| \\
&\le \sum_{d_1,d_2 \le N} |g(d_1)g(d_2)| \cdot \left| \sum_{n \le N/[d_1,d_2]} \chi(n) \right| \\
&\le \sum_{d_1,d_2 \le R} \left| \sum_{n \le N/[d_1,d_2]} \chi(n) \right|
\end{aligned}
$$

The inner sum can be estimated using either the Polya-Vinogradoff-inequality or Burgess estimates, leading to $|S| \le R^2\sqrt{q}\log q$ resp. $|S| \le c_{k,\epsilon} R^{2/k} N^{1-1/k} q^{(k+1)/(4k^2)+\epsilon}$, thus we obtain the third statement.

To prove the second statement, we begin as above to obtain the inequality

$$
\left| \sum_{n=1}^{\infty} f(n)\chi(n)e^{-\log^2(n/N)} \right| \le \sum_{d_1,d_2 \le R} \left| \sum_{n=1}^{\infty} \chi(n)e^{-\log^2([d_1,d_2]n/N)} \right|
$$

Write $d = [d_1, d_2]$. Using the Mellin-transform $\frac{1}{2\sqrt{\pi}i}\int_{(2)} x^{-s}e^{s^2/4}ds = e^{-\log^2 x}$, the inner sum can be expressed as

$$
\sum_{n=1}^{\infty} \chi(n)e^{-\log^2(dn/N)} = \frac{1}{2\sqrt{\pi}i}\int_{(2)} L(s,\chi)e^{s^2/4}(N/d)^s ds
$$

Now we shift the path of integration to the line $\Re s = -A$ with $A > 0$. Denote with χ_1 the primitive character inducing χ. Then we have

$$
L(s,\chi) = \prod_{p|q_2} \left(1 - \chi_1(p)p^{-s}\right) L(s,\chi_1).
$$

For $A > 2$, the first factor is $\ll q_2^A$, whereas the L-series can be estimated using the functional equation to be $\ll (q_1(|t|+2))^{A+1/2}$, hence the right hand side is $\ll_A q^{1/2}\left(\frac{N}{dq}\right)^{-A} \leq q^{1/2}\left(\frac{N}{R^2 q}\right)^{-A}$. Hence the whole sum can be bounded by $c(A)R^2 q^{1/2}\left(\frac{N}{R^2 q}\right)^{-A}$.

To prove Theorem 2, we follow the lines of the proof of the large sieve resp. the Halász-inequality, however, we apply Lemma 7 to a different euclidean space. Consider the subspace $V < l^\infty$ consisting of all bounded sequences (a_n), such that $a_n = 0$ whenever $f(n) = 0$, where f is defined as in Lemma 8. On this space define a product as $\langle (a_n), (b_n)\rangle := \sum_{n=1}^\infty f(n)e^{-\log^2(n/N)} a_n \overline{b_n}$. Now we apply Lemma 7 to this space and the set of vectors $\Phi = (\hat{a}_n)$, where $\hat{a}_p = a_p e^{\log^2 p/N}$, for prime numbers p in the range $R^2 < p \leq N$, and $\hat{a}_n = 0$ otherwise, and $v_i = (\hat{\chi}(n))$, where similary $\hat{\chi}(n) = \overline{\chi(n)}$, if $f(n) \neq 0$, and 0 otherwise. Now the inequality reads as

$$\sum_{q\leq Q}\sum_{\chi \pmod q}\left|\sum_{R^2<p\leq N} a_p\chi(p)\right|^2 \leq \max_\chi\left(\sum_{n=1}^\infty f(n)e^{-\log^2 n/N} + \sum_{\chi'\neq\chi}\left|\sum_{n\leq N} f(n)e^{-\log^2 n/N}\chi\overline{\chi'}(n)\right|\right) \times \sum_{p\leq N}|a_p|^2 e^{2\log^2(p/N)}$$

where the maximum is taken over all characters with moduli at most Q. From Lemma 8 it follows that the first term inside the brackets is $\ll \frac{N}{\log R}$, provided that $R < N^{1/3}$, say. For the second term, let χ be a charac-

ter $\pmod q$ and χ' a character $\pmod{q'}$. Then $\chi\overline{\chi'}$ is a character $\pmod{[q,q']}$. By Proposition 9, each term in the outer sum can be bounded by $c(A)R^2[q,q']^{1/2}\left(\frac{N}{R^2[q,q']}\right)^{-A}$, hence the whole sum is $\leq c(A)Q^3R^2\left(\frac{N}{R^2Q^2}\right)^{-A}$. Since by asumption $N > Q^{2+\epsilon}$, we can choose $R = Q^{\epsilon/4}$, $A = 6/\epsilon + 1$ to bound this by some constant depending only on ϵ. Thus we get the estimate

$$\sum_{q\leq Q}\sum_{\chi \pmod q}\left|\sum_{R^2<p\leq N} a_p\chi(p)\right|^2 \ll \left(\frac{N}{\epsilon\log N} + C_\epsilon\right)\sum_{p\leq N}|a_p|^2.$$

The range $n \leq R^2$ can be estimated using the usual large sieve inequality, which gives $(R^2+Q^2)\sum_{p\leq N}|a_p|^2$, which is negligible. Hence Theorem 2 is proven.

The proof of Theorem 3 is similar, but simpler. First, assume that all characters in $\mathcal{C}$ are characters to a single modulus q. We consider the vector space $V < \mathbb{C}^N$ consisting of sequences $(a_n)_{n=1}^N$ with $a_n = 0$ for all n with $f(n) = 0$ and the scalar product $\langle (a_n),(b_n)\rangle := \sum_{n\leq N} f(n)a_n\overline{b_n}$. Applying Lemma 7 as above, we obtain the estimate

$$\sum_{\chi\in\mathcal{C}}\left|\sum_{p\leq N} a_p\chi(p)\right|^2 \leq \left(\frac{N}{\log R} + R^2 + (|\mathcal{C}|-1)\Delta(R,N,q)\right)\sum_{R\leq p\leq N}|a_p|^2$$

where $\Delta(R,N,q)$ is the bound obtained by Proposition 9, i.e. $\Delta(R,N,q) \leq R^2\sqrt{q}\log q$, resp. $\Delta(R,N,q) < c_{k,\epsilon}q^{(k+1)/(4k^2)+\epsilon}N^{1-1/k}R^{2/k}$. The term R^2 can be neglected in comparison with $\Delta(R,N,q)$. This is obvious in the

first case. In the second case, we may assume that $\Delta(R, N, q) < N$, since otherwise Theorem 3 is an immediate consequence of the Cauchy-Schwarz-inequality. This implies $R < N^{1/2} q^{-(k+1)/(2k)}$, which in turn implies $R^2 < N^{1-1/k} q^{-1-1/k} < \Delta(R, N, q)$. Hence we obtain Theorem 3 for sets of characters belonging to a single modulus.

The proof for the case that the characters belong to different moduli is similar, note that $[q_1, q_2]$ is cubefree, if both q_1 and q_2 are cubefree.

In the range $Q^{2+\epsilon} \leq x < Q^{3+\epsilon}$, Corollary 4 follows from Theorem 2 by choosing $a_p = 1$ for all prime numbers $p \leq N$, whereas in the range $x > Q^{3+\epsilon}$ it follows from Theorem 3. Similarly we obtain corollary 5 from Theorem 3. We choose $\mathcal{C} = \{\chi_0, \chi, \overline{\chi}\}$ to obtain the estimate

$$|\pi(x)|^2 + 2|\pi(x, \chi)|^2 \leq \frac{x}{\log c_{k,\epsilon} x^{1/2} q^{(k+1)/(8k)+\epsilon}} \pi(x)$$

and choosing either $k = 3$ or $k \nearrow \infty$ we obtain the result by solving for $|\pi(x, \chi)|$.

To prove corollary 6, let $\mathcal{P}$ be the set of prime numbers p, such that there is some character χ of order d as described in the corollary. For every such p, choose such a character χ_1 together with all its powers, and denote

the set of all these character with $\mathcal{C}$. Let ζ be a d-th root of unity. We have

$$\sum_{\substack{\chi^d=\chi_0 \\ \chi\neq\chi_0}} |\pi(x,\chi)|^2 = d\sum_{a=1}^{d}\left|\#\{p\leq x|\chi_1(p)=\zeta^a\} - \frac{1}{d}\pi(x,\chi_0)\right|^2$$

Since by assumption, one of the terms on the right hand side is large, the right hand side is $\gg \frac{x^2}{d\log^2 x} \geq \frac{x}{D\log^2 x}$. Now we have $|\mathcal{C}| \leq D\cdot|\mathcal{P}|$, thus we get

$$|\mathcal{P}|\frac{x^2}{D\log^2 x} \ll \frac{x^2}{\log x\log R} + xDR^2|\mathcal{P}|Q\log Q$$

If $D^2Q\log Q < x^{1-\epsilon}$, we can choose $R = x^{\epsilon/4}$, and the second term on the right hand side is still of lesser order then the left hand side. With this choice the inequality can be simplified to $|\mathcal{P}| \ll_\epsilon D$.

References

[1] P. D. T. A. Elliott, *Subsequences of primes in residue classes to prime moduli*, in: Studies in pure mathematics to the Memory of P. Turán, P. Erdős (ed.), Akademia Kiado, Budapest, 1983, 157-164

[2] H. Halberstam, H.-E. Richert, *Sieve methods*, London Mathematical Society Monographs, No. 4, Academic Press, 1974

[3] H. L. Montgomery, *Topics in multplicative number theory*, Lecture Notes in Mathematics 227, Springer, 1971

[4] Y. Motohashi, *Large sieve extensions of the Brun-Titchmarsh theorem*, in: Studies in pure mathematics to the Memory of P. Turán, P. Erdős (ed.), Akademia Kiado, Budapest, 1983, 507-515

Jan-Christoph Puchta

Mathematical Institute

University of Oxford

24-29 St. Giles' Street

Oxford, OX1 3LB

United Kingdom

puchta@maths.ox.ac.uk